NIT-133-05

## IN THE CLAIMS

1-29. Canceled.

30. (Currently Amended) A cell components recovering apparatus comprising:

a substrate being disposed in a separation cell, wherein thea sample solution containing cells is supplied on a surface of the substrate, the substrate havingand a plurality of independent areas are formed on theits surface of the <del>oubstrate</del>;

capturing means for capturing each of the cells one by one separately on each of the plurality of independent areas; and

temperature control means for heating the surface of the substrate at one area of the plurality of independent areas to a predetermined temperature to destroy the cell captured at thesaid one area of the areas, to liberate cell components from the cell captured at the said one area. of the areas into the separation cell,

wherein, by introducing a washing solution into the separation cell, whereby the cells at the areas, except for the one area of the areas, remain on the areas, respectively,

NIT-133-05

the washing solution is recovered to recover the cell components liberated from the cell; and wherein, by changing-a position-of-the one area of the

areas, the washing-solution is recovered to recover the cell components liberated from the cell for each of the areas.

- 31. (Currently Amended) A cell component recovering apparatus according to claim 30, wherein the captured cell is a white blood cell.
- 32. (Previously Added) A cell component recovering apparatus according to claim 30, wherein the capturing means comprises means for applying an alternating field onto the surface of the substrate.
- 33. (Currently Amended) A cell components recovering apparatus according to claim 30, further comprising means for applying a DC field onto a surface of thesaid one area of the one area of the areas in a solution which contains no polynucleotide and has a pH value of 4 or lower, to attract nucleotide components to the surface of the one of the identified positions.

NIT-133-05

34. (Currently Amended) A cell components recovering apparatus comprising:

a substrate being disposed in a separation cell, wherein athe sample solution containing cells is supplied on a surface of the substrate, the substrate havingand a plurality of independent areas are-formed on theits surface of the substrate;

capturing means for capturing each of the cells one by one separately on each of the plurality of independent areas;

means for identifying athe positions of theone areas of said plurality of areas where athe cells to be destroyed areis present; and

temperature control means for heating the surface of the substrate at one of the identified positions said one area to a predetermined temperature to destroy the cell captured thereinat the area of the one of the identified positions, to liberate cell components from the cell captured thereinat the area of the identified positions into the separation cell.7

wherein, by introducing a washing solution into the separation cell, whereby the cells at the areas, except for the area at the one of the identified positions, remain on the areas, respectively, the washing solution is recovered to recover the cell components liberated from the cell; and

NIT-133-05

\_\_\_wherein, by changing a position of the identified positions, the weshing solution is recovered to recover the cell components liberated from the cell for each of the identified positions.

- 35. (Currently Amended) A cell component recovering apparatus according to claim 34, wherein the <u>captured</u> cell is a white blood cell.
- 36. (Previously Added) A cell component recovering apparatus according to claim 34, wherein the capturing means comprises means for applying an alternating field onto the surface of the substrate.
- 37. (Currently Amended) A cell component recovering apparatus according to claim 34, further comprising means for applying a DC field onto a surface of said one areathe one of the identified positions in a solution which contains no polynucleotide and has a pH value of 4 or lower, to attract nucleotide components to the surface of the one of the identified positions.
- 38. (New) A cell components recovering apparatus according to claim 30, wherein by introducing a washing

NIT-133-05

solution, the apparatus enables liberated cell components from the cell captured in said one area to be recovered while cells in areas other than said one area remain on the substrate,

and wherein said temperature control means individually heats different areas of said plurality of areas so as to enable recovery of cell components from all of said plurality of areas.

- 39. (New) A cell component recovering apparatus according to claim 38, wherein the captured cell is a white blood cell.
- 40. (New) A cell component recovering apparatus according to claim 38, wherein the capturing means comprises means for applying an alternating field onto the surface of the substrate.
- 41. (New) A cell components recovering apparatus according to claim 38, further comprising means for applying a DC field onto a surface of said one area in a solution which contains no polynucleotide and has a pH value of 4 or lower, to attract nucleotide components to the surface of the one of the identified positions.

NIT-133-05

42. (New) A cell components recovering apparatus according to claim 34, wherein by introducing a washing solution, the apparatus enables liberated cell components from the cell captured in said one area to be recovered while cells in areas other than said one area remain on the substrate,

and wherein said temperature control means individually heats different areas of said plurality of areas so as to enable recovery of cell components from all of said plurality of areas.

- 43. (New) A cell component recovering apparatus according to claim 42, wherein the captured cell is a white blood cell.
- 44. (New) A cell component recovering apparatus according to claim 42, wherein the capturing means comprises means for applying an alternating field onto the surface of the substrate.
- 45. (New) A cell components recovering apparatus according to claim 42, further comprising means for applying a DC field onto a surface of said one area in a solution which contains no polynucleotide and has a pH value of 4 or lower, to attract nucleotide components to the surface of the one of the identified positions.